

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
14 November 2002 (14.11.2002)

PCT

(10) International Publication Number  
**WO 02/090367 A1**(51) International Patent Classification<sup>7</sup>: **C07F 9/38**

European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(21) International Application Number: **PCT/TR02/00018**(22) International Filing Date: **8 May 2002 (08.05.2002)**(25) Filing Language: **English**(26) Publication Language: **English**(30) Priority Data:  
2001/01250 10 May 2001 (10.05.2001) **TR**(71) Applicant (for all designated States except US): **EOS ECZACIBASI OZGUN KIMYASAL URUNLER SANAYI VE TICARET A.S.** [TR/TR]; Buyukdere Caddesi, Ali Kaya Sokak No. 7, Levent, 80640 Istanbul (TR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DABAK, Kadir** [TR/TR]; Buyukdere Caddesi, Ali Kaya Sokak No. 7, Levent, 80640 Istanbul (TR). **OZARSLAN, A., Evren** [TR/TR]; Buyukdere Caddesi, Ali Kaya Sokak No. 7, Levent, 80640 Istanbul (TR). **SAHBAZ, Filiz** [TR/TR]; Buyukdere Caddesi, Ali Kaya Sokak No. 7, Levent, 80640 Istanbul (TR). **ASLAN, Tuncer** [TR/TR]; Buyukdere Caddesi, Ali Kaya Sokak No. 7, Levent, 80640 Istanbul (TR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- of inventorship (Rule 4.17(iv)) for US only

**Published:**

- with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **PROCESS FOR THE PREPARATION OF 4-AMINO-1-HYDROXYBUTYLIDENE-1, 1-BIPHOSPHONIC ACID**

(57) **Abstract:** This invention is related with the preparation of 4-amino-1-hydroxybutylidene-1,1-biphosphonic acid or salts thereof. The reaction of 4-aminobutyric acid with phosphorous acid and phosphorous trichloride in the presence of aralkyl or alkyl ethoxylates or triglycerides such as plant or animal oils or their derivative; and recovering of 4-amino-1-hydroxybutylidene-1,1-biphosphonic acid or salts thereof are described. The main feature of the invention is in the use of the above defined non-ionic emulgators in the phosphorylation reaction.

**WO 02/090367 A1**

PROCESS FOR THE PRODUCTION OF  
4-AMINO-1-HYDROXYBUTYLIDENE-1, 1-BIPHOSPHONIC ACID

Process for the preparation of 4-amino-1-hydroxybutylidene-1,1-biphosphonic acid or salts thereof

This invention is related with the preparation of 4-amino-1-hydroxybutylidene-1,1-biphosphonic acid or salts thereof. The reaction of 4-aminobutyric acid with  
5 phosphorous acid and phosphorus trichloride in the presence of aralkyl or alkyl ethoxylates or triglycerides such as plant or animal oils or their derivatives and recovering of 4-amino-1-hydroxybutylidene-1,1-biphosphonic acid or salts thereof are described.

4-Amino-1-hydroxybutylidene-1,1-biphosphonic acid monosodium salt trihydrate is  
10 used for treatment or prevent of diseases involving bone disorders, such as hypercalcemia of malignanch, Paget's disease and osteoporosis.

4-Amino-1-hydroxybutylidene-1,1-biphosphonic acid or salts thereof are prepared basically by the reaction of 4-aminobutyric acid with a mixture of phosphorous acid and one of the three phosphorus chlorides; phosphorous trichloride, phosphorous  
15 oxychloride or phosphorous pentachloride and then quenching the reaction mixture with water followed by heating to hydrolyse the phosphorous intermediates.

Several patented methods can be found in the literature for the preparation of  $\omega$ -amino-1-hydroxyalkylidene-1,1-bisphosphonic acids and especially for 4-amino-1-hydroxybutylidene-1,1-biphosphonic acid and salts thereof. In U.S. Patent 4,407,761  
20 (Blom et al.) the preparation of 4-amino-1-hydroxybutylidene-1,1-biphosphonic acid besides other bisphosphonic acids are described. When using this procedure, a semisolid sticky non-stirrable mass develops which prevents smooth heat transfer. The described process might be suitable for laboratory preparations, however for industrial production it is not acceptable. In U.S. Patent 4,705,651 (Staibano, G.), a  
25 similar procedure is followed with different molar ratios and although some improvements were achieved, it is still unsuitable for industrial scale up.

Kieczkowski et al. (In U.S. Patents 4,922,007; 5,019,651 and J. Org. Chem. 1995, 60, 8310-8312) reported that the solidification problem has been solved. Methanesulfonic acid was used to solubilize the reaction components and keep them  
30 fluid throughout. By the use of methanesulfonic acid, the fluidity problems were solved however another serious safety problem surfaced. A reaction between methanesulfonic acid and phosphorus trichloride is exothermic and at certain point becomes uncontrollable.

U.S. Patent 5,908,959 (Kubela et al.) also describes the preparation of 4-amino-1-  
5 hydroxybutylidene-1,1-biphosphonic acid or salts thereof. The reaction is carried out in poly(alkylene glycol) as a diluent, which solubilizes the reaction components, however still when the reaction mixture is decomposed with water, an agitation problem occurs. The viscous reaction mixture must be transferred into the water. To facilitate this, viscosity problem is solved by the addition of toluene. When using  
10 toluene, a safety problem arises and also an additional separation step is needed.

In the present invention; by the use of aralkyl or alkyl ethoxylates or triglycerides such as plant or animal oils or their derivatives as emulgators, the solidification and the safety problems are solved in a cheaper, safer and easily accessible way without any need of an additional solvent. These emulgators solubilize the reaction  
15 components and do not react with the reactants to cause any uncontrolled reactions. It has been found that 4-amino-1-hydroxybutylidene-1,1-biphosphonic acid or its salts can be obtained in a safe and high yielded way without an additional purification step.

Aralkyl or alkyl ethoxylates or their derivatives are used very often in the textile,  
20 leather and metal industries as emulgators. Triglycerides such as plant or animal oils are also used very often in the food and lubricant industries. These emulgators are easily accessible, readily available and non-expensive.

The reaction of 4-aminobutyric acid with phosphorous acid and phosphorus trichloride in the presence of one of these emulgators at a suitable temperature such  
25 as between about 40 °C and about 150 °C; and hydrolysing the phosphorous

intermediates by heating the reaction mixture in the presence of water and recovering of 4-amino-1-hydroxybutylidene-1,1-biphosphonic acid or salts thereof are described.

Aralkyl or alkyl ethoxylates or their derivatives may be selected from the general  
30 formula of  $R-X-O-(CH_2CH_2O)_n-H$ ; and triglycerides might be selected from the general formula of  $(RCO_2CH_2)_2CHCOOR$  wherein R represents branched or non-branched alkyl groups which contain 1 to 20 carbon atoms, X represents phenyl or naphthyl or  $-CH_2-$  groups and; n is a number of between 1 and about 30.

The main feature of the invention is in the use of the above defined emulsifiers or  
5 derivatives thereof in the phosphorylation reaction. These compounds keep the mixture in homogenous form and can be separated easily from the product at the end of the reaction and can be reused. The hydrolysis of the formed phosphorous intermediates can be completed in the same reaction mixture and if desired, by adjusting the pH to about 4.3, the sodium salt of the said biphosphonic acid can be  
10 directly obtained and isolated in a pure form.

The 4-aminobutyric acid and the phosphorous acid are suspended in one of the mentioned emulgators and reacted with phosphorus trichloride at a suitable temperature for example between about 40 °C and about 150 °C, preferably at about 70 °C. The phosphorylation reaction is completed in about 3 hours at this  
15 temperature. The preferred ratio of the amino acid to phosphorous acid and to phosphorous trichloride is about 1:1:2.

As example of aralkyl or alkyl ethoxylates or their derivatives, which can be applied, are nonylphenol with 4 mol, 6 mol or 10 mol ethoxylate; and alkyl ethoxylates such as lauryl alcohol of different ethoxylate numbers. And as example  
20 of triglycerids are sunflower oil, olive oil and corn oil, which are glyceride of oleic acid, palmitic acid, linoleic acid, stearic acid, myristic acid, behenic acid and arachidic acid in different ratios.

The reaction can be shown schematically as follows:

PLEASE INSERT SCHEME HERE

Wherein R represents branched or non-branched alkyl groups which contain 1 to 20  
25 carbon atoms, X represents phenyl or naphthyl or  $-\text{CH}_2-$  groups and; n is a number of  
between 1 and about 30.

The following examples are introduced the practical procedures and the results  
without any limitations in any subject.

**EXAMPLE 1**

30 Preparation of 4-amino-1-hydroxybutylidene-1,1-biphosphonic Acid Monosodium  
Salt Trihydrate in Nonylphenol Ethoxylate 4 Mol (NP4) (or in Nonylphenol  
Ethoxylate 6 Mol)

A 2-L flask charged with 500 mL of NP 4 (or Nonylphenol Ethoxylate 6 Mol), 75.4  
g (0.73 mol) of 4-aminobutyric acid and 60 g (0.73 mol) of phosphorous acid in  
5 room temperature. The system was connected to a caustic scrubber and flushed with  
nitrogen. After 15 minutes of stirring, 132 mL of phosphorous trichloride was added  
by dropwise addition over a period of 30 minutes. Then the reaction mixture was  
stirred at 70 °C for 4 hours. After 4 hours, the mixture was cooled to 20 °C, then 300  
mL of water was added by dropwise addition over a period of 30 minutes. After  
10 completion of the addition of the water, the reaction mixture was heated at 105 °C  
for 4 hours, then cooled to 20 °C. The stirring was discontinued to allow the layers  
separate, the lower aqueous layer was separated and the pH of this solution was  
adjusted to 4.3 with 50% NaOH. After stirring for 13 hours, 50 mL of acetone was  
added and stirred for 1 hr, then the crystalline product was collected by filtration,

15 washed with 100 mL of ice cold water and 100 mL of acetone and dried at room temperature. The yield of 4-amino-1-hydroxybutylidene-1,1-biphosphonic acid monosodium salt trihydrate was 136.5 g (57.4 %). The analysis confirmed the identity of the product and the absence of impurities.

<sup>32</sup>P-NMR, <sup>1</sup>H-NMR and <sup>13</sup>C-NMR analyses were recorded on a Varian Mercury 300  
20 MHz instrument.

<sup>32</sup>P-NMR (D<sub>2</sub>O), 18.794 (s); <sup>13</sup>C-NMR (D<sub>2</sub>O), 30.68 (t), 40.07(t), 73.59 (s); <sup>1</sup>H-NMR (D<sub>2</sub>O), 1.84 (4H, m), 2.87 (2H, m).

## EXAMPLE 2

Preparation of 4-amino-1-hydroxybutylidene-1,1-biphosphonic Acid Monosodium  
25 Salt Trihydrate in Nonylphenol Ethoxylate 10 Mol

Instead of NP4, NP10 was used as a solvent and a procedure was followed as described in Example 1 until 4-hour hydrolysis at about 110 °C and cooling to room temperature. Then pH of all the reaction mixture (because two phases were not formed at this point as described in example 1.) was adjusted to 4.3 and then two  
30 phases were formed. The lower phase was separated, after stirring for 13 hours, 50 mL of acetone was added and stirred for 1 hr. Precipitated crystalline product was collected by filtration, washed with 100 mL of ice cold water, 100 mL of acetone and dried at room temperature. The analysis confirmed the identity of the product and the absence of impurities. The yield of 4-amino-1-hydroxybutylidene-1,1-biphosphonic acid monosodium salt trihydrate was 58%.

## EXAMPLE 3

Preparation of 4-amino-1-hydroxybutylidene-1,1-biphosphonic Acid in  
Nonylphenol Ethoxylate 4 Mol (or in Nonylphenol Ethoxylate 6 Mol)

A procedure was followed as described in Example 1 until 4-hour hydrolysis at  
10 about 110 °C and cooling to room temperature. Then stirring was discontinued to allow the layers separate, the lower aqueous layer was separated and 500 mL of

acetone was added and stirred. The product firstly was separated as in oil form then was crystallised after 10 minutes stirring. The crystalline product was collected by filtration, washed with 100 mL of ice cold water and 100 mL of acetone and dried at  
15 room temperature. The yield of 4-amino-1-hydroxybutylidene-1,1-biphosphonic Acid was 60 %. 1 L of acetone was added to the upper layer and the yield was increased from 60 to 65% by the filtration of precipitated product however since this second precept was not as pure as the first one, it needs an additional crystallisation. The analysis confirmed the identity of the product and the absence of impurities.

20

#### EXAMPLE 4

Preparation of 4-amino-1-hydroxybutylidene-1,1-biphosphonic Acid in Nonylphenol Ethoxylate 10 Mol

Instead of NP4, NP10 was used as a solvent and a procedure was followed as described in Example 1 until 4-hour hydrolysis at about 110 °C and cooling to room  
25 temperature. Then 1,5 L of acetone was added and stirred, the product firstly was separated as in oil form then was crystallised after 10 minutes stirring. The crystalline product was collected by filtration, washed with 250 mL of ice cold water and 250 mL of acetone and dried at room temperature. The yield of 4-amino-1-hydroxybutylidene-1,1-biphosphonic Acid was 60 %. The analysis confirmed the  
30 identity of the product and the absence of impurities.

#### EXAMPLE 5

Preparation of 4-amino-1-hydroxybutylidene-1,1-biphosphonic Acid Monosodium Salt Trihydrate in Lauryl alcohol Ethoxylate 6 Mol

Instead of NP4, lauryl alcohol Ethoxylate 6 Mol was used and an experiment was  
5 carried out same as it was described in Example 1. The yield of 4-amino-1-hydroxybutylidene-1,1-biphosphonic Acid Monosodium Salt Trihydrate was (59 %). The analysis confirmed the identity of the product and the absence of impurities.

**EXAMPLE 6**

Preparation of 4-Amino-1-hydroxybutylidene-1, 1-bisphosphonic acid monosodium  
10 salt trihydrate by using sunflower oil.

A 3 L flask was equipped with a mechanical stirrer, thermometer, condenser, and an addition funnel. The system was connected to a caustic scrubber and flushed with nitrogen. The flask was charged with 500 mL of sunflower oil. The temperature was brought to 75 °C. At this temperature 100 g (0.97 mol) of 4-aminobutyric acid and  
15 79.5 g (0.97 mol) of phosphorous acid were added. The mixture was stirred for 15 minutes. Phosphorous trichloride 130 mL (1.45 mol) was added to this solution in 20 minutes by keeping the internal temperature between 70 to 75 °C. The mixture was stirred at this temperature for 3 hours and then 500 mL of water was added in portions. The mixture is stirred for 10 minutes, and transferred into a separatory  
20 funnel and separated. The aqueous phase was heated at 105 °C for 6 hours. The pH of the solution was brought to 4.3 by adding 50% NaOH. The solution was concentrated half of its volume (250 mL) and stirred for 12 hours at 25 °C. The product was collected by filtration washed with 25 mL of cold water, air dried at 25 °C to give 135 g of 4-Amino-1-hydroxybutylidene-1, 1-bisphosphonic acid  
25 monosodium salt trihydrate as a white solid in 43% yield. After completion of the reaction, the structure of the sunflower oil was checked by <sup>1</sup>H-NMR and <sup>13</sup>C-NMR and no change was observed. This result showed that the oil can be reused.

**EXAMPLE 7**

30 Preparation of 4-Amino-1-hydroxybutylidene-1, 1-bisphosphonic acid monosodium salt trihydrate by using recovered sunflower oil.

A 1 L flask was equipped with a mechanical stirrer, thermometer, condenser, and an addition funnel. The system was connected to a caustic scrubber and flushed with nitrogen. The flask was charged with 100 mL of recovered sunflower oil obtained  
5 from example 1. The temperature was brought to 75 °C. At this temperature, 20 g (0.19 mol) of 4-aminobutyric acid and 15.9 g (0.19 mol) of phosphorous acid were



added. The mixture was stirred for 15 minutes. Phosphorous trichloride 26 mL (0.29 mol) was added to this solution in 10 minutes by keeping the internal temperature between 70 °C to 75 °C. The mixture was stirred at this temperature for 3 hours and  
10 then 100 mL of water was added in portions. The mixture is stirred for 5 minutes, and transferred into a separatory funnel. The phases were separated. The aqueous phase was taken and stirred at 105 °C for 6 hours. The pH of the solution was brought to 4.3 by adding 50% NaOH. The solution was stirred for 12 hours 25 °C. The product was collected by filtration, washed with 25 mL of cold water, air dried  
15 at 25 °C to give 26.3 g of 4-Amino-1-hydroxybutylidene-1, 1-bisphosphonic acid monosodium salt trihydrate as a white solid in 42% yield.

### EXAMPLE 8

Preparation of 4-Amino-1-hydroxybutylidene-1, 1-bisphosphonic acid monosodium salt trihydrate by using sunflower oil without separation of the phases before  
20 hydrolysis.

A 500 mL flask was equipped with a mechanical stirrer, thermometer, condenser, and an addition funnel. The system was connected to a caustic scrubber and flushed with nitrogen. The flask was charged with 50 mL of sunflower oil. The temperature was brought to 75 °C. At this temperature 10 g (0.097 mol) of 4-aminobutyric acid  
25 and 7.95 g (0.097 mol) of phosphorous acid were added. The mixture was stirred for 15 minutes. Phosphorous trichloride 13 mL (0.145 mol) was added to this solution in 5 minutes by keeping the internal temperature between 70 to 75 °C. The mixture was stirred for 3 hours at this temperature and then 50 mL of water was added. The two phases system was stirred at 105 °C for 6 hours and then transferred into a  
30 separatory funnel. The aqueous phase was taken. The pH of the solution was brought to 4.3 by adding 50% NaOH. Acetone 25 mL was added to the solution and stirred for 12 hours 25 °C. The product was collected by filtration washed with 25 mL of cold water, air dried at 25 °C to give 13.2 g of 4-Amino-1-hydroxybutylidene-1, 1-bisphosphonic acid monosodium salt trihydrate as a white solid in 43% yield.  
5 The structure of the oil used in this reaction was checked by <sup>1</sup>H-NMR, the spectra showed some hydrolysis of the triglycerides under reflux condition.

**EXAMPLE 9**

Preparation of 4-Amino-1-hydroxybutylidene-1, 1-bisphosphonic acid monosodium salt trihydrate by using olive oil.


- 10 Instead of sunflower oil, olive oil was used as a solvent. The reaction was carried out with a 10 g scale following the procedure described in example 1. After filtration, 9 g of 4-Amino-1-hydroxybutylidene-1, 1-bisphosphonic acid monosodium salt trihydrate was obtained as a white solid in 36% yield.

## CLAIMS

What is claimed is:

1. A process is provided for the preparation of 4-amino-1-hydroxybutylidene 1,1-biphosphonic acid or salts thereof which comprises of:
  - (a) reacting 4-aminobutyric acid with phosphorous acid and phosphorus trichloride  
5 in the presence of aralkyl or alkyl ethoxylates or their derivatives which might be selected from the general formula of  $R-X-O-(CH_2CH_2O)_n-H$  or, triglycerides such as plant or animal oils or their derivatives which might be selected from the general formula of  $(RCO_2CH_2)_2CHCOOR$  wherein R represents branched or non-branched alkyl or alkenyl groups which contain 1 to 20 carbon atoms, X represents phenyl or  
10 naphthyl or  $-CH_2-$  and; n is a number of between 1 and about 30.
  - (b) recovering said 4-amino-1-hydroxybutylidene-1,1-biphosphonic acid or salts thereof by the hydrolysis reaction of phosphorous intermediates.
2. The process of Claim 1 when the said reaction is carried out in the presence of such ethoxylates or triglycerides wherein  $R = C_4-C_{19}$  alkyl or alkenyl,  $X =$  phenyl or  
15  $-CH_2-$ , and  $n = 4-30$ .
3. The process of Claim 2 wherein said reaction is carried out at temperature of from  $40^\circ C$  to  $150^\circ C$ .
4. The process of Claim 2 when the said reaction is conducted in the presence of such ethoxylate wherein  $R = C_9$ ,  $X =$  phenyl and  $n = 4-10$ . Such compounds are  
20 nonylphenol with 4 mol, 6 mol or 10 mol ethoxylate.
5. The process of Claim 2 when the said reaction is conducted in the presence of such ethoxylate wherein  $R = C_{11}$ ,  $X = CH_2$  and  $n = 6$ . Such compound is lauryl alcohol with 6 mol ethoxylate.
6. The process of Claim 2 when the said reaction is conducted in the presence of  
25 such triglyceride wherein  $R = C_{12}-C_{20}$  which can contain one or more double bonds. Such compounds are sunflower oil, olive oil and corn oil.

7. The process of claim 4, 5 and 6 wherein the temperature is about 70 °C.
8. The process of Claim 4,5 and 6 wherein 4-amino-1-hydroxybutylidene-1,1-biphosphonic acid or its salts thereof is recovered.



SCHEME

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/TR 02/00018

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 C07F9/38

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 922 007 A (GERARD R. KIECZYKOWSKI) 1 May 1990 (1990-05-01) cited in the application the whole document	1-8
A	WO 98 34940 A (APOTEX INC.) 13 August 1998 (1998-08-13) cited in the application the whole document	1-8

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

30 July 2002

Date of mailing of the international search report

08/08/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Beslier, L

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No  
PCT/TR 02/00018

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 4922007	A	01-05-1990	AT 129713 T	15-11-1995
			AU 625704 B2	16-07-1992
			AU 5701990 A	13-12-1990
			CA 2018477 A1	09-12-1990
			DE 69023280 D1	07-12-1995
			DE 69023280 T2	20-06-1996
			DK 402152 T3	04-12-1995
			EP 0402152 A2	12-12-1990
			ES 2080116 T3	01-02-1996
			FI 93219 B	30-11-1994
			GR 3018379 T3	31-03-1996
			HK 69596 A	26-04-1996
			HU 9500204 A3	28-08-1995
			IE 69564 B1	02-10-1996
			IL 94612 A	30-03-1995
			JP 7048391 A	21-02-1995
			JP 1931325 C	12-05-1995
			JP 3101684 A	26-04-1991
			JP 6062651 B	17-08-1994
			KR 137455 B1	01-05-1998
			LV 11472 A	20-08-1996
			LV 11472 B	20-12-1996
			NO 902559 A ,B,	10-12-1990
			NO 941726 A ,B,	10-12-1990
			NZ 233972 A	26-05-1992
			PT 94306 A ,B	08-02-1991
			ZA 9004446 A	24-06-1992
WO 9834940	A	13-08-1998	CA 2197267 A1	11-08-1998
			AU 728164 B2	04-01-2001
			AU 5977298 A	26-08-1998
			BR 9807568 A	21-03-2000
			WO 9834940 A1	13-08-1998
			EP 0971938 A1	19-01-2000
			PL 335053 A1	27-03-2000
			US 5908959 A	01-06-1999